050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_negative_max_area_40C
Pearson r=-0.351 (p=0.0531) | Spearman ρ=-0.370 (p=0.0405) | Kendall τ=-0.276 (p=0.0329)
Pearson log1p(activity) r=-0.332 (p=0.0681)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.35, p=0.328)

Global statistics\nn = 31\nR² = 0.123\nRMSE = 69.22 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.33 (p=0.466)
  Spearman ρ=0.17 (p=0.711)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.35 (p=0.328)
  Spearman ρ=0.27 (p=0.448)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.50 (p=0.173)
  Spearman ρ=-0.35 (p=0.349)