050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_positive_total_area_40C
Pearson r=0.548 (p=0.00143) | Spearman ρ=0.530 (p=0.00218) | Kendall τ=0.384 (p=0.00285)
Pearson log1p(activity) r=0.459 (p=0.00935)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.40, p=0.255)

Global statistics\nn = 31\nR² = 0.300\nRMSE = 61.85 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.07 (p=0.882)
  Spearman ρ=0.20 (p=0.67)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.40 (p=0.255)
  Spearman ρ=-0.32 (p=0.372)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.05 (p=0.891)
  Spearman ρ=0.16 (p=0.68)