069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=0.005 (p=0.978) | Spearman ρ=-0.194 (p=0.295) | Kendall τ=-0.116 (p=0.366)
Pearson log1p(activity) r=-0.077 (p=0.681)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.24, p=0.496)

Global statistics\nn = 31\nR² = 0.000\nRMSE = 73.91 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=0.50 (p=0.253)
Spearman ρ=0.45 (p=0.31)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.24 (p=0.496)
Spearman ρ=-0.23 (p=0.519)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.17 (p=0.663)
Spearman ρ=-0.16 (p=0.68)