069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_negative_max_area_40C
Pearson r=-0.109 (p=0.56) | Spearman ρ=-0.031 (p=0.867) | Kendall τ=-0.018 (p=0.891)
Pearson log1p(activity) r=0.014 (p=0.939)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.49, p=0.151)

Global statistics\nn = 31\nR² = 0.012\nRMSE = 73.47 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.36 (p=0.43)
  Spearman ρ=-0.44 (p=0.328)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.49 (p=0.151)
  Spearman ρ=0.38 (p=0.273)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.09 (p=0.827)
  Spearman ρ=0.16 (p=0.68)