069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_positive_max_area_40C
Pearson r=-0.158 (p=0.396) | Spearman ρ=-0.199 (p=0.284) | Kendall τ=-0.139 (p=0.289)
Pearson log1p(activity) r=-0.142 (p=0.446)




Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.32, p=0.36)

Global statistics\nn = 31\nR² = 0.025\nRMSE = 72.98 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.01 (p=0.983)
  Spearman ρ=-0.30 (p=0.513)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.32 (p=0.36)
  Spearman ρ=-0.38 (p=0.275)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.01 (p=0.975)
  Spearman ρ=-0.16 (p=0.687)