089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.214 (p=0.247) | Spearman ρ=-0.203 (p=0.273) | Kendall τ=-0.148 (p=0.258)
Pearson log1p(activity) r=-0.202 (p=0.277)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.10, p=0.787)

Global statistics\nn = 31\nR² = 0.046\nRMSE = 72.20 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.19 (p=0.687)
  Spearman ρ=0.06 (p=0.907)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.10 (p=0.787)
  Spearman ρ=0.04 (p=0.913)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.31 (p=0.424)
  Spearman ρ=-0.31 (p=0.416)