089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.119 (p=0.524) | Spearman ρ=-0.235 (p=0.202) | Kendall τ=-0.172 (p=0.183)
Pearson log1p(activity) r=-0.161 (p=0.388)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.10, p=0.78)

Global statistics\nn = 31\nR² = 0.014\nRMSE = 73.38 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.19 (p=0.677)
  Spearman ρ=-0.13 (p=0.788)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.10 (p=0.78)
  Spearman ρ=-0.14 (p=0.705)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.02 (p=0.956)
  Spearman ρ=-0.08 (p=0.846)