108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.319 (p=0.08) | Spearman ρ=-0.348 (p=0.0549) | Kendall τ=-0.236 (p=0.0701)
Pearson log1p(activity) r=-0.380 (p=0.0352)




Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.11, p=0.768)

Global statistics\nn = 31\nR² = 0.102\nRMSE = 70.04 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.40 (p=0.378)
  Spearman ρ=-0.48 (p=0.274)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.11 (p=0.768)
  Spearman ρ=-0.04 (p=0.906)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.05 (p=0.898)
  Spearman ρ=0.09 (p=0.812)