108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_total_area_40C
Pearson r=0.507 (p=0.00361) | Spearman ρ=0.531 (p=0.00213) | Kendall τ=0.369 (p=0.00417)
Pearson log1p(activity) r=0.489 (p=0.00525)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.19, p=0.601)

Global statistics\nn = 31\nR² = 0.257\nRMSE = 63.71 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.27 (p=0.565)
  Spearman ρ=-0.28 (p=0.54)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.19 (p=0.601)
  Spearman ρ=-0.15 (p=0.687)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.41 (p=0.267)
  Spearman ρ=0.40 (p=0.283)