127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_positive_total_area_40C
Pearson r=0.556 (p=0.00117) | Spearman ρ=0.583 (p=5.74e-04) | Kendall τ=0.458 (p=3.66e-04)
Pearson log1p(activity) r=0.585 (p=5.43e-04)




Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.09, p=0.803)

Global statistics\nn = 31\nR² = 0.309\nRMSE = 61.44 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.49 (p=0.268)
  Spearman ρ=-0.27 (p=0.558)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.09 (p=0.803)
  Spearman ρ=0.19 (p=0.606)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.56 (p=0.114)
  Spearman ρ=0.47 (p=0.199)