146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.215 (p=0.246) | Spearman ρ=-0.075 (p=0.69) | Kendall τ=-0.049 (p=0.707)
Pearson log1p(activity) r=-0.215 (p=0.245)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.64, p=0.0472)

Global statistics\nn = 31\nR² = 0.046\nRMSE = 72.19 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.57 (p=0.178)
  Spearman ρ=0.32 (p=0.487)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.64 (p=0.0472)
  Spearman ρ=0.88 (p=8.01e-04)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.42 (p=0.263)
  Spearman ρ=-0.30 (p=0.44)