146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.137 (p=0.463) | Spearman ρ=-0.250 (p=0.174) | Kendall τ=-0.200 (p=0.124)
Pearson log1p(activity) r=-0.131 (p=0.484)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.20, p=0.585)

Global statistics\nn = 31\nR² = 0.019\nRMSE = 73.21 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.18 (p=0.692)
  Spearman ρ=-0.13 (p=0.786)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.20 (p=0.585)
  Spearman ρ=-0.38 (p=0.28)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.05 (p=0.895)
  Spearman ρ=-0.24 (p=0.54)