001 | orientation selected from FORMULA global_score
FORMULA_global_score_50C
Pearson r=-0.437 (p=0.014) | Spearman ρ=-0.246 (p=0.182) | Kendall τ=-0.190 (p=0.138)
Pearson log1p(activity) r=-0.291 (p=0.113)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.01, p=0.989)

Global statistics\nn = 31\nR² = 0.191\nRMSE = 80.25 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.02 (p=0.969)
  Spearman ρ=0.07 (p=0.878)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.51 (p=0.128)
  Spearman ρ=0.72 (p=0.018)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.01 (p=0.989)
  Spearman ρ=0.01 (p=0.983)