001 | orientation selected from FORMULA global_score
FORMULA_negative_max_area_50C
Pearson r=-0.299 (p=0.103) | Spearman ρ=-0.096 (p=0.606) | Kendall τ=-0.086 (p=0.515)
Pearson log1p(activity) r=-0.144 (p=0.441)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.05, p=0.907)

Global statistics\nn = 31\nR² = 0.089\nRMSE = 85.14 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.02 (p=0.963)
  Spearman ρ=-0.04 (p=0.938)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.11 (p=0.761)
  Spearman ρ=0.24 (p=0.511)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.05 (p=0.907)
  Spearman ρ=0.09 (p=0.811)