001 | orientation selected from FORMULA global_score
FORMULA_positive_total_area_50C
Pearson r=0.407 (p=0.023) | Spearman ρ=0.360 (p=0.0469) | Kendall τ=0.259 (p=0.0442)
Pearson log1p(activity) r=0.390 (p=0.03)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.42, p=0.266)

Global statistics\nn = 31\nR² = 0.166\nRMSE = 81.49 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.17 (p=0.708)
  Spearman ρ=-0.05 (p=0.908)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.07 (p=0.849)
  Spearman ρ=-0.02 (p=0.953)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.42 (p=0.266)
  Spearman ρ=0.28 (p=0.47)