012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_positive_total_area_50C
Pearson r=0.544 (p=0.00155) | Spearman ρ=0.391 (p=0.0296) | Kendall τ=0.274 (p=0.0331)
Pearson log1p(activity) r=0.401 (p=0.0253)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.27, p=0.484)

Global statistics\nn = 31\nR² = 0.296\nRMSE = 74.85 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.47 (p=0.283)
  Spearman ρ=0.38 (p=0.403)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.36 (p=0.308)
  Spearman ρ=-0.25 (p=0.486)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.27 (p=0.484)
  Spearman ρ=-0.22 (p=0.578)