031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=0.322 (p=0.0769) | Spearman ρ=-0.046 (p=0.804) | Kendall τ=-0.038 (p=0.771)
Pearson log1p(activity) r=0.176 (p=0.344)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.23, p=0.554)

Global statistics\nn = 31\nR² = 0.104\nRMSE = 84.45 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.49 (p=0.259)
  Spearman ρ=0.50 (p=0.253)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.03 (p=0.941)
  Spearman ρ=0.26 (p=0.464)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.23 (p=0.554)
  Spearman ρ=0.36 (p=0.335)