069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_global_score_50C
Pearson r=0.021 (p=0.909) | Spearman ρ=-0.164 (p=0.377) | Kendall τ=-0.098 (p=0.443)
Pearson log1p(activity) r=-0.036 (p=0.847)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.32, p=0.394)

Global statistics\nn = 31\nR² = 0.000\nRMSE = 89.19 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.41 (p=0.358)
  Spearman ρ=0.40 (p=0.379)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.11 (p=0.764)
  Spearman ρ=0.01 (p=0.973)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.32 (p=0.394)
  Spearman ρ=-0.43 (p=0.25)