069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=-0.042 (p=0.823) | Spearman ρ=-0.102 (p=0.585) | Kendall τ=-0.058 (p=0.656)
Pearson log1p(activity) r=-0.077 (p=0.682)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.08, p=0.831)

Global statistics\nn = 31\nR² = 0.002\nRMSE = 89.14 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.06 (p=0.906)
  Spearman ρ=0.01 (p=0.985)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.24 (p=0.499)
  Spearman ρ=-0.17 (p=0.646)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.08 (p=0.831)
  Spearman ρ=-0.07 (p=0.854)