069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_positive_total_area_50C
Pearson r=0.550 (p=0.00134) | Spearman ρ=0.423 (p=0.0177) | Kendall τ=0.317 (p=0.014)
Pearson log1p(activity) r=0.417 (p=0.0196)




Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.26, p=0.501)

Global statistics\nn = 31\nR² = 0.303\nRMSE = 74.50 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.57 (p=0.179)
  Spearman ρ=0.59 (p=0.159)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.05 (p=0.892)
  Spearman ρ=-0.21 (p=0.566)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.26 (p=0.501)
  Spearman ρ=-0.29 (p=0.442)