078 | orientation selected from PEARSON global_score
PEARSON_negative_max_area_50C
Pearson r=-0.209 (p=0.259) | Spearman ρ=-0.088 (p=0.638) | Kendall τ=-0.055 (p=0.68)
Pearson log1p(activity) r=-0.135 (p=0.469)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.38, p=0.313)

Global statistics\nn = 31\nR² = 0.044\nRMSE = 87.25 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.05 (p=0.924)
  Spearman ρ=0.03 (p=0.954)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.11 (p=0.764)
  Spearman ρ=0.31 (p=0.381)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.38 (p=0.313)
  Spearman ρ=0.48 (p=0.188)