
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_global_score_50C
Pearson r=0.189 (p=0.307) | Spearman ρ=0.283 (p=0.123) | Kendall τ=0.190 (p=0.138)
Pearson log1p(activity) r=0.337 (p=0.0634)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG3 Cutinase / PETase-like regression (r=0.38, p=0.313)
Global statistics\nn = 31\nR² = 0.036\nRMSE = 87.60 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.02 (p=0.964)
Spearman ρ=-0.07 (p=0.878)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.02 (p=0.959)
Spearman ρ=-0.23 (p=0.521)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.38 (p=0.313)
Spearman ρ=0.41 (p=0.271)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
GatA
nc_NylC
HotPETase
NfpolyA
UMGSP1
FsC
ThcCut1
LCC_ICCG
UMGSP3
ThcCut2
Tfcut2
HiC
ReU
TtA
Ooh
NylB
AoC
Trypsin
MGS0156
ph_NylC
Subtilisin
Cut190
MsmeCA
Papain
chymotrypsin