089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.025 (p=0.892) | Spearman ρ=-0.134 (p=0.472) | Kendall τ=-0.071 (p=0.585)
Pearson log1p(activity) r=-0.095 (p=0.612)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.00, p=0.997)

Global statistics\nn = 31\nR² = 0.001\nRMSE = 89.19 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.20 (p=0.672)
  Spearman ρ=0.14 (p=0.758)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.46 (p=0.178)
  Spearman ρ=-0.13 (p=0.712)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.00 (p=0.997)
  Spearman ρ=0.11 (p=0.78)