108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_negative_total_area_50C
Pearson r=0.514 (p=0.00312) | Spearman ρ=0.388 (p=0.0311) | Kendall τ=0.272 (p=0.0345)
Pearson log1p(activity) r=0.412 (p=0.0214)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.29, p=0.456)

Global statistics\nn = 31\nR² = 0.264\nRMSE = 76.55 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.12 (p=0.804)
  Spearman ρ=0.07 (p=0.877)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.42 (p=0.232)
  Spearman ρ=-0.44 (p=0.206)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.29 (p=0.456)
  Spearman ρ=0.19 (p=0.618)