108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_max_area_50C
Pearson r=-0.221 (p=0.232) | Spearman ρ=-0.163 (p=0.381) | Kendall τ=-0.133 (p=0.305)
Pearson log1p(activity) r=-0.255 (p=0.166)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.03, p=0.934)

Global statistics\nn = 31\nR² = 0.049\nRMSE = 87.01 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.40 (p=0.377)
  Spearman ρ=-0.43 (p=0.339)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.02 (p=0.946)
  Spearman ρ=0.15 (p=0.671)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.03 (p=0.934)
  Spearman ρ=0.24 (p=0.542)