108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_total_area_50C
Pearson r=0.425 (p=0.0172) | Spearman ρ=0.421 (p=0.0182) | Kendall τ=0.302 (p=0.0186)
Pearson log1p(activity) r=0.415 (p=0.0201)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.55, p=0.127)

Global statistics\nn = 31\nR² = 0.180\nRMSE = 80.76 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=-0.26 (p=0.571)
Spearman ρ=-0.18 (p=0.696)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.19 (p=0.6)
Spearman ρ=-0.20 (p=0.578)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.55 (p=0.127)
Spearman ρ=0.61 (p=0.0826)