127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.279 (p=0.129) | Spearman ρ=-0.200 (p=0.28) | Kendall τ=-0.146 (p=0.26)
Pearson log1p(activity) r=-0.220 (p=0.234)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.17, p=0.669)

Global statistics\nn = 31\nR² = 0.078\nRMSE = 85.68 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.14 (p=0.762)
  Spearman ρ=0.41 (p=0.357)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.03 (p=0.938)
  Spearman ρ=0.08 (p=0.834)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.17 (p=0.669)
  Spearman ρ=-0.06 (p=0.88)