127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_positive_total_area_50C
Pearson r=0.495 (p=0.0046) | Spearman ρ=0.518 (p=0.00283) | Kendall τ=0.392 (p=0.00228)
Pearson log1p(activity) r=0.534 (p=0.00198)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.70, p=0.0367)

Global statistics\nn = 31\nR² = 0.245\nRMSE = 77.50 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.51 (p=0.239)
  Spearman ρ=-0.36 (p=0.427)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.08 (p=0.819)
  Spearman ρ=0.16 (p=0.649)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.70 (p=0.0367)
  Spearman ρ=0.69 (p=0.04)