146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_total_area_50C
Pearson r=0.411 (p=0.0217) | Spearman ρ=0.315 (p=0.0845) | Kendall τ=0.239 (p=0.0631)
Pearson log1p(activity) r=0.355 (p=0.0501)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.77, p=0.0151)

Global statistics\nn = 31\nR² = 0.169\nRMSE = 81.34 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.72 (p=0.0681)
  Spearman ρ=-0.74 (p=0.0579)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.28 (p=0.442)
  Spearman ρ=-0.40 (p=0.253)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.77 (p=0.0151)
  Spearman ρ=0.62 (p=0.0738)