146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_positive_total_area_50C
Pearson r=0.524 (p=0.00247) | Spearman ρ=0.530 (p=0.00218) | Kendall τ=0.381 (p=0.00301)
Pearson log1p(activity) r=0.504 (p=0.00387)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.56, p=0.121)

Global statistics\nn = 31\nR² = 0.275\nRMSE = 75.98 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=0.60 (p=0.151)
Spearman ρ=0.45 (p=0.306)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.07 (p=0.842)
Spearman ρ=0.16 (p=0.65)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.56 (p=0.121)
Spearman ρ=0.60 (p=0.0899)