
001 | orientation selected from FORMULA global_score
FORMULA_global_score_50C
Pearson r=-0.437 (p=0.014) | Spearman ρ=-0.246 (p=0.182) | Kendall τ=-0.190 (p=0.138)
Pearson log1p(activity) r=-0.291 (p=0.113)
PA6 activity (µM)
SURFMATCH global score
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
UMGSP1
FsC
HotPETase
NfpolyA
UMGSP3
HiC
LCC_ICCG
ThcCut1
ph_NylC
TtA
ReU
MsmegA
NylB
Ooh
Tfcut2
Trypsin
Papain
Cut190
MGS0156
ThcCut2
AoC
chymotrypsin
Subtilisin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.191\nRMSE = 80.25 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.02 (p=0.969)
Spearman ρ=0.07 (p=0.878)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.51 (p=0.128)
Spearman ρ=0.72 (p=0.018)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.01 (p=0.989)
Spearman ρ=0.01 (p=0.983)