
001 | orientation selected from FORMULA global_score
FORMULA_negative_total_area_50C
Pearson r=0.507 (p=0.00357) | Spearman ρ=0.354 (p=0.0508) | Kendall τ=0.229 (p=0.076)
Pearson log1p(activity) r=0.396 (p=0.0274)
PA6 activity (µM)
SURFMATCH score_negative_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.258\nRMSE = 76.87 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.40 (p=0.374)
Spearman ρ=0.38 (p=0.403)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.04 (p=0.913)
Spearman ρ=-0.06 (p=0.867)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.49 (p=0.176)
Spearman ρ=-0.40 (p=0.282)
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
ThcCut1
LCC_ICCG
HiC
Tfcut2
Cut190
UMGSP1
NfpolyA
UMGSP3
ThcCut2
TtA
ph_NylC
MGS0156
AoC
ReU
MsmegA
NylB
Ooh
Trypsin
Papain
chymotrypsin
Subtilisin