
001 | orientation selected from FORMULA global_score
FORMULA_positive_max_area_50C
Pearson r=-0.143 (p=0.442) | Spearman ρ=-0.261 (p=0.156) | Kendall τ=-0.188 (p=0.146)
Pearson log1p(activity) r=-0.140 (p=0.452)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
UMGSP1
NfpolyA
FsC
LCC_ICCG
AoC
ThcCut1
UMGSP3
HiC
TtA
Trypsin
ph_NylC
ThcCut2
Tfcut2
ReU
Cut190
MGS0156
NylB
Subtilisin
MsmegA
Papain
Ocn
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.021\nRMSE = 88.30 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.19 (p=0.681)
Spearman ρ=-0.16 (p=0.728)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.30 (p=0.402)
Spearman ρ=-0.08 (p=0.834)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.22 (p=0.561)
Spearman ρ=-0.14 (p=0.714)