
001 | orientation selected from FORMULA global_score
FORMULA_positive_total_area_50C
Pearson r=0.407 (p=0.023) | Spearman ρ=0.360 (p=0.0469) | Kendall τ=0.259 (p=0.0442)
Pearson log1p(activity) r=0.390 (p=0.03)
PA6 activity (µM)
SURFMATCH score_positive_total_area
0
50
100
150
200
250
300
300
320
340
360
380
400
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
ThcCut1
LCC_ICCG
HiC
AoC
ThcCut2
Trypsin
Cut190
Papain
Tfcut2
Subtilisin
Ooh
MGS0156
Chymotrypsin
NylB
UMGSP1
NfpolyA
UMGSP3
TtA
ph_NylC
MsmegA
ReU
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.166\nRMSE = 81.49 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.17 (p=0.708)
Spearman ρ=-0.05 (p=0.908)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.07 (p=0.849)
Spearman ρ=-0.02 (p=0.953)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.42 (p=0.266)
Spearman ρ=0.28 (p=0.47)