
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.228 (p=0.217) | Spearman ρ=-0.099 (p=0.596) | Kendall τ=-0.061 (p=0.644)
Pearson log1p(activity) r=-0.120 (p=0.52)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.052\nRMSE = 86.86 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.46 (p=0.301)
Spearman ρ=-0.19 (p=0.682)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.12 (p=0.75)
Spearman ρ=-0.26 (p=0.46)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.17 (p=0.659)
Spearman ρ=0.12 (p=0.761)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
ThcCut1
AoC
UMGSP1
UMGSP3
FsC
NfpolyA
ReU
HiC
TtA
Tfcut2
LCC_ICCG
ThcCut2
ph_NylC
MGS0156
Trypsin
NylB
MsmegA
Ooh
Subtilisin
Papain
Cut190
chymotrypsin