
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_negative_total_area_50C
Pearson r=0.493 (p=0.00481) | Spearman ρ=0.329 (p=0.071) | Kendall τ=0.234 (p=0.0682)
Pearson log1p(activity) r=0.370 (p=0.0404)
PA6 activity (µM)
SURFMATCH score_negative_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.243\nRMSE = 77.61 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.56 (p=0.187)
Spearman ρ=0.58 (p=0.175)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.20 (p=0.585)
Spearman ρ=-0.01 (p=0.987)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.00 (p=0.992)
Spearman ρ=0.00 (p=1)
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
ThcCut1
UMGSP1
NfpolyA
LCC_ICCG
UMGSP3
HiC
ph_NylC
Tfcut2
ThcCut2
TtA
MGS0156
Papain
Cut190
ReU
Trypsin
NylB
MsmegA
AoC
Ooh1
chymotrypsin
Subtilisin