
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=-0.007 (p=0.97) | Spearman ρ=-0.105 (p=0.575) | Kendall τ=-0.079 (p=0.539)
Pearson log1p(activity) r=-0.063 (p=0.736)
PA6 activity (µM)
SURFMATCH score_positive_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.000\nRMSE = 89.21 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.12 (p=0.805)
Spearman ρ=0.11 (p=0.818)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.47 (p=0.171)
Spearman ρ=0.36 (p=0.306)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.35 (p=0.353)
Spearman ρ=-0.38 (p=0.311)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
NfpolyA
UMGSP1
FsC
LCC_ICCG
ThcCut1
UMGSP3
Tfcut2
TtA
NylB
ph_NylC
ThcCut2
HiC
ReU
MsmegA
Ooh
Cut190
MGS0156
AoC
Trypsin
Subtilisin
chymotrypsin
Papain