
031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_global_score_50C
Pearson r=-0.294 (p=0.109) | Spearman ρ=-0.145 (p=0.438) | Kendall τ=-0.111 (p=0.385)
Pearson log1p(activity) r=-0.126 (p=0.499)
PA6 activity (µM)
SURFMATCH global score
0
50
100
150
200
250
300
0.00
0.05
0.10
0.15
0.20
0.25
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
UMGSP1
NfpolyA
ThcCut1
FsC
HotPETase
UMGSP3
LCC_ICCG
Tfcut2
ThcCut2
HiC
TtA
ph_NylC
Trypsin
ReU
AoC
NylB
Cut190
Ooh
MsmegA
Papain
chymotrypsin
Subtilisin
MGS0156
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.086\nRMSE = 85.28 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.24 (p=0.599)
Spearman ρ=0.02 (p=0.969)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.31 (p=0.378)
Spearman ρ=0.26 (p=0.476)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.62 (p=0.0734)
Spearman ρ=0.55 (p=0.121)