031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_negative_total_area_50C
Pearson r=0.551 (p=0.00133) | Spearman ρ=0.388 (p=0.0309) | Kendall τ=0.236 (p=0.0657)
Pearson log1p(activity) r=0.428 (p=0.0163)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression

Global statistics\nn = 31\nR² = 0.303\nRMSE = 74.47 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=0.87 (p=0.0109)
Spearman ρ=0.81 (p=0.0269)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.13 (p=0.713)
Spearman ρ=-0.16 (p=0.65)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.57 (p=0.113)
Spearman ρ=-0.51 (p=0.158)