
031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=0.322 (p=0.0769) | Spearman ρ=-0.046 (p=0.804) | Kendall τ=-0.038 (p=0.771)
Pearson log1p(activity) r=0.176 (p=0.344)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
250
300
20
30
40
50
60
70
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
UMGSP1
NfpolyA
FsC
ThcCut1
LCC_ICCG
Tfcut2
UMGSP3
ThcCut2
HiC
ph_NylC
ReU
TtA
MsmegA
NylB
MGS0156
Trypsin
Subtilisin
Ooh
Cut190
AoC
Papain
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.104\nRMSE = 84.45 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.49 (p=0.259)
Spearman ρ=0.50 (p=0.253)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.03 (p=0.941)
Spearman ρ=0.26 (p=0.464)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.23 (p=0.554)
Spearman ρ=0.36 (p=0.335)