
050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_global_score_50C
Pearson r=0.089 (p=0.634) | Spearman ρ=0.255 (p=0.166) | Kendall τ=0.168 (p=0.189)
Pearson log1p(activity) r=0.216 (p=0.243)
PA6 activity (µM)
SURFMATCH global score
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
ThcCut1
HiC
LCC_ICCG
UMGSP3
NfpolyA
AoC
Tfcut2
Papain
TtA
Trypsin
MGS0156
ReU
Cut190
ph_NylC
Ooh
NylB
ThcCut2
Subtilisin
MsmegA
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.008\nRMSE = 88.86 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.08 (p=0.858)
Spearman ρ=-0.38 (p=0.403)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.26 (p=0.467)
Spearman ρ=0.21 (p=0.567)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.13 (p=0.732)
Spearman ρ=0.02 (p=0.966)