
050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.268 (p=0.146) | Spearman ρ=-0.243 (p=0.187) | Kendall τ=-0.192 (p=0.138)
Pearson log1p(activity) r=-0.244 (p=0.186)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.072\nRMSE = 85.96 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.42 (p=0.344)
Spearman ρ=0.45 (p=0.317)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.08 (p=0.816)
Spearman ρ=0.31 (p=0.377)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.46 (p=0.208)
Spearman ρ=-0.36 (p=0.339)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GalA
HotPETase
FsC
NfpolyA
UMGSP1
ThcCut1
UMGSP3
LCC_ICCG
HiC
TTcut2
ThcCut2
TtA
ph_NylC
ReU
NylB
MGS0156
Trypsin
AoC
MsmegA
Cut190
Subtilisin
Papain
chymotrypsin