
050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_positive_max_area_50C
Pearson r=-0.065 (p=0.726) | Spearman ρ=0.076 (p=0.683) | Kendall τ=0.038 (p=0.771)
Pearson log1p(activity) r=0.041 (p=0.825)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
UMGSP1
HotPETase
FsC
NfpolyA
ThcCut1
LCC_ICCG
UMGSP3
HiC
Tfcut2
TtA
Ooh
NylB
ReU
MsmegA
chymotrypsin
Trypsin
ThcCut2
Cut190
Subtilisin
ph_NylC
MGS0156
AoC
Papain
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.004\nRMSE = 89.02 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.03 (p=0.953)
Spearman ρ=0.20 (p=0.67)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.49 (p=0.15)
Spearman ρ=0.45 (p=0.193)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.08 (p=0.84)
Spearman ρ=0.27 (p=0.475)