050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_positive_total_area_50C
Pearson r=0.532 (p=0.00206) | Spearman ρ=0.480 (p=0.00633) | Kendall τ=0.339 (p=0.00824)
Pearson log1p(activity) r=0.480 (p=0.00626)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression

Global statistics\nn = 31\nR² = 0.283\nRMSE = 75.53 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.13 (p=0.785)
  Spearman ρ=-0.11 (p=0.818)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.01 (p=0.985)
  Spearman ρ=-0.17 (p=0.638)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.13 (p=0.736)
  Spearman ρ=0.21 (p=0.587)