
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_global_score_50C
Pearson r=0.021 (p=0.909) | Spearman ρ=-0.164 (p=0.377) | Kendall τ=-0.098 (p=0.443)
Pearson log1p(activity) r=-0.036 (p=0.847)
PA6 activity (µM)
SURFMATCH global score
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
ThcCut1
NfpolyA
LCC_ICCG
UMGSP3
HiC
Tfcut2
TtA
ReU
ph_NylC
Trypsin
ThcCut2
Cut190
chymotrypsin
Ooh
AoC
NylB
MGS0156
MsmegA
Papain
Subtilisin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.000\nRMSE = 89.19 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.41 (p=0.358)
Spearman ρ=0.40 (p=0.379)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.11 (p=0.764)
Spearman ρ=0.01 (p=0.973)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.32 (p=0.394)
Spearman ρ=-0.43 (p=0.25)