
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_negative_max_area_50C
Pearson r=-0.135 (p=0.468) | Spearman ρ=0.093 (p=0.62) | Kendall τ=0.060 (p=0.645)
Pearson log1p(activity) r=0.075 (p=0.687)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.018\nRMSE = 88.39 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.45 (p=0.316)
Spearman ρ=-0.50 (p=0.253)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.15 (p=0.68)
Spearman ρ=0.26 (p=0.466)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.18 (p=0.65)
Spearman ρ=0.33 (p=0.389)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
NfpolyA
FsC
UMGSP1
ThcCut1
UMGSP3
LCC_ICCG
Tfcut2
ThcCut2
ph_NylC
HiC
MsmegA
TtA
Papain
ReU
NylB
Ooh
Trypsin
MGS0156
AoC
Cut190
Subtilisin
chymotrypsin