
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_positive_total_area_50C
Pearson r=0.550 (p=0.00134) | Spearman ρ=0.423 (p=0.0177) | Kendall τ=0.317 (p=0.014)
Pearson log1p(activity) r=0.417 (p=0.0196)
PA6 activity (µM)
SURFMATCH score_positive_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.303\nRMSE = 74.50 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.57 (p=0.179)
Spearman ρ=0.59 (p=0.159)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.05 (p=0.892)
Spearman ρ=-0.21 (p=0.566)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.26 (p=0.501)
Spearman ρ=-0.29 (p=0.442)
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
NfpolyA
UMGSP3
ThcCut1
LCC_ICCG
HiC
AoC
chymotrypsin
ThcCut2
Trypsin
Subtilisin
Tfcut2
MGS0156
TtA
ReU
Ooh
Cut190
Papain
NylB
MsmegA
ph_NylC