
078 | orientation selected from PEARSON global_score
PEARSON_negative_max_area_50C
Pearson r=-0.209 (p=0.259) | Spearman ρ=-0.088 (p=0.638) | Kendall τ=-0.055 (p=0.68)
Pearson log1p(activity) r=-0.135 (p=0.469)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.044\nRMSE = 87.25 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.05 (p=0.924)
Spearman ρ=0.03 (p=0.954)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.11 (p=0.764)
Spearman ρ=0.31 (p=0.381)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.38 (p=0.313)
Spearman ρ=0.48 (p=0.188)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
GatA
nc_NylC
HotPETase
FsC
ThcCut1
UMGSP1
NfpolyA
LCC_ICCG
UMGSP3
HiC
ph_NylC
Tfcut2
TtA
NylB
MsmegA
Ooh
Cut190
ThcCut2
ReU
MGS0156
AoC
chymotrypsin
Trypsin
Papain
Subtilisin