
078 | orientation selected from PEARSON global_score
PEARSON_positive_total_area_50C
Pearson r=0.406 (p=0.0233) | Spearman ρ=0.434 (p=0.0146) | Kendall τ=0.287 (p=0.0256)
Pearson log1p(activity) r=0.433 (p=0.0149)
PA6 activity (µM)
SURFMATCH score_positive_total_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.165\nRMSE = 81.52 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.16 (p=0.734)
Spearman ρ=-0.05 (p=0.908)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.13 (p=0.728)
Spearman ρ=-0.18 (p=0.614)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.58 (p=0.103)
Spearman ρ=0.57 (p=0.108)
NylCk
NylCp2_TS
NylCp2
NylCa
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
NfpolyA
UMGSP3
TtA
ph_NylC
ReU
MsmegA
HiC
NylB
Papain
Ooh
Cut190
MGS0156
Trypsin
chymotrypsin
Subtilisin
LCC_ICCG
ThcCut1
Tfcut2
ThcCut2
AoC