
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_negative_max_area_50C
Pearson r=-0.190 (p=0.305) | Spearman ρ=-0.174 (p=0.348) | Kendall τ=-0.108 (p=0.411)
Pearson log1p(activity) r=-0.237 (p=0.199)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.036\nRMSE = 87.59 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.07 (p=0.883)
Spearman ρ=0.10 (p=0.83)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.17 (p=0.638)
Spearman ρ=0.17 (p=0.63)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.38 (p=0.311)
Spearman ρ=-0.18 (p=0.635)
NylCk
NylCp2_TS
NylCa
NylCp2
tt_NylC
UMGSP2
nc_NylC
GatA
HotPETase
FsC
UMGSP1
NfpolyA
LCC_ICCG
ThcCut1
UMGSP3
HiC
ph_NylC
Tfcut2
ThcCut2
TtA
NylB
ReU
Trypsin
Subtilisin
Papain
AoC
MsmegA
Cut190
MGS0156
Ooh
chymotrypsin